#### European Environment Agency European Topic Centre on Biological Diversity



### Aquila fasciatus

Annex I Yes-CTC International action plan SAP

Bonelli's Eagle, *Aquila fasciatus*, is a species of day-flying bird of prey found in woodland and forest and unvegetated or sparsely vegetated land ecosystems. It breeds across the Southern and South-Eastern Europe. It tends to live in warm, mountainous regions, nesting on cliff edges and sometimes on trees. Typically, vegetation in these areas is dominated by scrub, but it also inhabits more densely covered areas and almost completely bare areas (www.arkive.org)

Aquila fasciatus has a breeding population size of 1000-1200 pairs and a breeding range size of 282000 square kilometres in the EU27. The breeding population trend in the EU27 is Stable in the short term and Decreasing in the long term.

The EU population status of *Aquila fasciatus* was assessed as Near Threatened, because the species comes close to meeting the IUCN Red List criteria at the EU27 scale.


## Aquila fasciatus


Report under the Article 12 of the Birds Directive

# Assessment of status at the European level

Breeding population size	Breeding population trend		_ Range _	Breeding range trend		Winter	Winter population trend		Population
	Short term	Long term	area	Short term	Long term	population size	Short term	Long term	status
1000 - 1200 p	0	-	282000						Near Threatened

See the endnotes for more information<sup>i</sup>







## Trends at the Member State level

MS/Ter. % in EU27 po		Breeding	Breeding population trend		Range	Breeding range trend		Winter	Winter population trend	
		population size	Short term	Long term	area	Short term	Long term	population size	Short term	Long term
BG	0.3	1 - 3 p	X	X	400	0	0			
CY	2.6	40 - 50 p	+	+	5000	+	+			
ES	73.9	741 - 763 p	0	-	202787	0	-			
FR	3.3	28 - 31 p	+	-	9100	+	-			
GR										
IT	7.3	25 - 30 p	X	_	17100	-	+			
PT	12.7	128 - 150 p	+	+	48500	+	+			

See the endnotes for more information<sup>ii</sup>





Short-term winter population trend was not reported for this species.

### Main pressures and threats reported by Member States

For the bird species triggering SPA classification Member States were asked to report the 20 most important pressures and threats using an agreed hierarchical list which can be found on the Article 12 Reference Portal (http://bd.eionet.europa.eu/activities/Reporting/Article\_12/reference\_portal). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. The table below only contains information from Member States, where a species triggers SPA classification. Pressures and threats were ranked in three classes 'high, medium and low importance', the table below only shows pressures and threats classed as 'high', for some species there were less than ten pressures and threats reported as highly important.

#### Ten most frequently reported 'highly important' pressures and threats

Code	Activity	Frequency
D02	Utility and service lines/pipelines	18
F03	Hunting and collection of terrestrial wild animals	18
J03	Other changes to ecosystems	18
G01	Outdoor sports, leisure and recreational activities	12
A02	Modification of cultivation practices	6
A04	Grazing by livestock	6
C01	Mining and quarrying	6
C03	Production of renewable energy (abiotic)	6
D01	Roads, railroads and paths	6
G05	Other human intrusions and disturbances	6

### Proportion of population covered by the Natura 2000 network

For the bird species triggering SPA classification Member States were asked to report the size of a species population occurring within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size.

#### Percentage of coverage by Natura 2000 sites

MS/territory	season	SPA trigger	% coverage
BG	breeding	NO	
CY	breeding	YES	43.3
ES	breeding	YES	0.8
FR	breeding	YES	89.93
IT	breeding	YES	51.64

#### Aquila fasciatus

Report under the Article 12 of the Birds Directive

MS/territory	season	SPA trigger	% coverage
PT	breeding	YES	41.5

See the endnotes for more informationiii

### Most frequently reported conservation measures

For the bird species triggering SPA classification Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 12 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

#### Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.1	Establish protected areas/sites	27
2.1	Maintaining grasslands and other open habitats	9
3.0	Other forestry-related measures	9
3.2	Adapt forest management	9
6.0	Other spatial measures	9
6.3	Legal protection of habitats and species	9
7.1	Regulation/ Management of hunting and taking	9
7.4	Specific single species or species group management measures	9
8.2	Specific management of traffic and energy transport systems	9

This information is derived from the Member State national reports submitted to the European Commission under Article 12 of the Birds Directive in 2013 and covering the period 2008-2012. More detailed information, including the MS reports, is available at: http://bd.eionet.europa.eu/article12/summary?period=1&subject=A707.

#### Aquila fasciatus

Report under the Article 12 of the Birds Directive

<sup>1</sup> Assessment of status at the European level: The EU assessments of birds population status was made by the European Red List of Birds Consortium (under contract with the European Commission)

The EU27 population trends were assessed using these categories: '+' Increasing, '0' Stable, 'F' Fluctuating, '-' Decreasing, 'xu' Uncertain and 'x' Unknown. The breeding population size is estimated in majority of the cases as 'p' number of pairs. Alternative population units used are: 'males' number of males, 'i' number of individuals, 'cmales' number of calling males and 'bfem' number of breeding females. The winter population size is estimated as number of individuals.

<sup>ii</sup>Species trends at the Member State level: The percentage of the EU27 species population occurring in the Member States (% in EU27) is calculated based on the population size reported by the Member States.

iiiPercentage of coverage by Natura 2000 sites: In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (\*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece and for non-Annex I species in the Czech Republic.