# Report under the Article 12 of the Birds Directive Period 2008-2012

### **European Environment Agency** European Topic Centre on Biological Diversity



### Calidris canutus

Annex I No International action plan No

Red Knot, *Calidris canutus*, is a species of wader found in unvegetated or sparsely vegetated land and marine inlet and transitional water ecosystems. It breeds in Greenland, but winters mainly in coastal areas of western Europe. This species is a full long-distance migrant that utilises few stopover sites or staging areas. The species breeds on dry upland tundra including weathered sandstone ridges, upland areas with scattered willows (*Salix spp.*), *Dryas spp.* and poppy, moist marshy slopes and flats in foothills, well-drained slopes hummocked with *Dryas spp.* and upland glacial gravel close to streams or ponds. Outside of the breeding season the species is strictly coastal, frequenting tidal mudflats or sandflats, sandy beaches of sheltered coasts, rocky shelves, bays, lagoons and harbours, occasionally also oceanic beaches and saltmarshes (European Red List 2015).

Calidris canutus has a winter population size of 504000-565000 individuals in the EU27. The winter population trend in the EU27 is Increasing in the short term and Increasing in the long term.

The EU population status of *Calidris canutus* was assessed as Secure, because the species does not meet any of the IUCN Red List criteria for threatened or Near Threatened, or the criteria for Depleted or Declining (the EU27 population or range has not declined by 20% or more since 1980).

## Assessment of status at the European level

Breeding population size	Breeding population trend		_ Range _	Breeding range trend		_ Winter	Winter population trend		_Population
	Short term	Long term	area	Short term	Long term	population size	Short term	Long term	status
						504000 - 565000 i	+	+	Secure

See the endnotes for more information<sup>i</sup>


Distribution map not available.


### Trends at the Member State level

MS/Ter. % in	Breeding population size	Breeding population trend		_ Range _	Breeding range trend		_ Winter	Winter population trend	
MS/Ter. EU27		Short term	Long term	area	Short term	Long term	population size	Short term	Long term
BE							15 - 126 i	X	X
BG							0 - 6 i	X	X
DE							37000 - 37000 i	+	F
DK							27000 - 27000 i	F	F
ES							329 - 541 i	+	-
FR							36000 - 58000 i	+	+
IE							22120 - 22120 i	+	X
NL							51321 - 89006 i	F	0
PT							1122 - 1122 i	0	0
UK							330000 - 330000 i	+	+

See the endnotes for more information  $^{\mathrm{ii}}$ 

Short-term breeding population trend was not reported for this species.





### Main pressures and threats reported by Member States

For the bird species triggering SPA classification Member States were asked to report the 20 most important pressures and threats using an agreed hierarchical list which can be found on the Article 12 Reference Portal (http://bd.eionet.europa.eu/activities/Reporting/Article\_12/reference\_portal). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. The table below only contains information from Member States, where a species triggers SPA classification. Pressures and threats were ranked in three classes 'high, medium and low importance', the table below only shows pressures and threats classed as 'high', for some species there were less than ten pressures and threats reported as highly important.

### Ten most frequently reported 'highly important' pressures and threats

Code	Activity	Frequency
F02	Fishing and harvesting aquatic resources	67
F06	Other hunting, fishing and collection activities	33

### Proportion of population covered by the Natura 2000 network

For the bird species triggering SPA classification Member States were asked to report the size of a species population occurring within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size.

#### Percentage of coverage by Natura 2000 sites

MS/territory	season	SPA trigger	% coverage
BE	winter	NO	
BG	winter	YES	100
DE	winter	YES	100
DK	winter	YES	97.96
ES	winter	YES	23.47
FR	winter	YES	92.9
IE	winter	YES	85.15
NL	winter	NO	
PT	winter	NO	
UK	winter	YES	86.26

See the endnotes for more information iii

### Most frequently reported conservation measures

For the bird species triggering SPA classification Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be

### Calidris canutus

Report under the Article 12 of the Birds Directive

found on the Article 12 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

#### Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.1	Establish protected areas/sites	47
6.3	Legal protection of habitats and species	35
9.2	Regulating/Managing exploitation of natural resources on sea	12
7.2	Regulation/ Management of fishery in limnic systems	6

This information is derived from the Member State national reports submitted to the European Commission under Article 12 of the Birds Directive in 2013 and covering the period 2008-2012. More detailed information, including the MS reports, is available at: <a href="http://bd.eionet.europa.eu/article12/summary?period=1&subject=A143">http://bd.eionet.europa.eu/article12/summary?period=1&subject=A143</a>.

### Calidris canutus

Report under the Article 12 of the Birds Directive

<sup>1</sup> Assessment of status at the European level: The EU assessments of birds population status was made by the European Red List of Birds Consortium (under contract with the European Commission)

The EU27 population trends were assessed using these categories: '+' Increasing, '0' Stable, 'F' Fluctuating, '-' Decreasing, 'xu' Uncertain and 'x' Unknown. The breeding population size is estimated in majority of the cases as 'p' number of pairs. Alternative population units used are: 'males' number of males, 'i' number of individuals, 'cmales' number of calling males and 'bfem' number of breeding females. The winter population size is estimated as number of individuals.

<sup>ii</sup>Species trends at the Member State level: The percentage of the EU27 species population occurring in the Member States (% in EU27) is calculated based on the population size reported by the Member States.

iiiPercentage of coverage by Natura 2000 sites: In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (\*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece and for non-Annex I species in the Czech Republic.