



Chlidonias niger

Annex I	Yes
International action plan	No

Black Tern, *Chlidonias niger*, is a species of seabird found in wetland and river and lake ecosystems.

Chlidonias niger has a breeding population size of 11 100-20600 pairs and a breeding range size of 125000 square kilometres in the EU27. The breeding population trend in the EU27 is Decreasing in the short term and Decreasing in the long term.



The EU population status of *Chlidonias niger* was assessed as Declining, because EU27 population or range declined by at least 20% since 1980, with continuing decline since 2001.

Assessment of status at the European level

Breeding population size	Breeding population trend		Range area	Breeding range trend		Winter population size	Winter population trend		Population status
	Short term	Long term		Short term	Long term		Short term	Long term	
11100 - 20600 p	-	-	125000						Declining

See the endnotes for more informationⁱ

Chlidonias niger
Report under the Article 12 of the Birds Directive





Trends at the Member State level

MS/Ter.	% in EU27	Breeding population size	Breeding population trend		Range area	Breeding range trend		Winter population size	Winter population trend	
			Short term	Long term		Short term	Long term		Short term	Long term
BG	0.4	25 - 52 p	-	-	300	-	-			
CZ										
DE	14.4	900 - 1100 p	0	0	10981	0	0			
DK	1.1	48 - 58 p	-	-	406	0	-			
EE	12.8	1000 - 2000 p	-	-	9200	-	-			
ES	0.7	0 - 40 p	F	-	526	F	-			
FI	2.4	15 - 25 p	-	0	2900	x	-			
FR	2.7	95 - 270 p	-	F	2600	0	-			
GR										
HU	9.9	100 - 1400 p	F	-	5358	F	F			
IT	1.8	120 - 120 p	x	F	1200	-	-			
LT	19.3	3000 - 6000 p	+	0	33500	0	0			
LV	18.8	2118 - 3842 p	x	+	38180	x	0			
NL	6.4	1150 - 1450 p	0	-	7967	-	-			
PL		2000 - 3000 p	-	-		x	x			
RO	2.6	300 - 800 p	x	x	3000	x	x			
SE	5.3	250 - 300 p	0	+	7900	0	x			
SK	1.5	0 - 10 p	-	-	1300	-	-			

See the endnotes for more informationⁱⁱ

Chlidonias niger

Report under the Article 12 of the Birds Directive



Short-term winter population trend was not reported for this species.

Main pressures and threats reported by Member States

For the bird species triggering SPA classification Member States were asked to report the 20 most important pressures and threats using an agreed hierarchical list which can be found on the Article 12 Reference Portal (http://bd.eionet.europa.eu/activities/Reporting/Article_12/reference_portal). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. The table below only contains information from Member States, where a species triggers SPA classification. Pressures and threats were ranked in three classes 'high, medium and low importance', the table below only shows pressures and threats classed as 'high', for some species there were less than ten pressures and threats reported as highly important.

Ten most frequently reported 'highly important' pressures and threats

Code	Activity	Frequency
J02	Changes in water bodies conditions	30
A02	Modification of cultivation practices	13
J03	Other changes to ecosystems	13
H01	Pollution to surface waters	10
A01	Agricultural cultivation	3
A07	Use of 'pesticides' in agriculture	3
A09	Irrigation in agriculture	3
F01	Marine and freshwater aquaculture	3
F02	Fishing and harvesting aquatic resources	3
F03	Hunting and collection of terrestrial wild animals	3

Proportion of population covered by the Natura 2000 network

For the bird species triggering SPA classification Member States were asked to report the size of a species population occurring within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size.

Percentage of coverage by Natura 2000 sites

MS/territory	season	SPA trigger	% coverage
BG	breeding	YES	7.84
DE	breeding	YES	92.92
DK	breeding	YES	92.38
EE	breeding	YES	47.43
ES	breeding	YES	x

MS/territory	season	SPA trigger	% coverage
FI	breeding	YES	36.51
FR	breeding	YES	83.77
HU	breeding	YES	82.11
IT	breeding	YES	91.29
LT	breeding	YES	7.35
LV	breeding	YES	33.19
NL	breeding	YES	52.08
PL	breeding	YES	x
RO	breeding	YES	89.86
SE	breeding	YES	85.73
SK	breeding	YES	100

See the endnotes for more informationⁱⁱⁱ

Most frequently reported conservation measures

For the bird species triggering SPA classification Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 12 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.1	Establish protected areas/sites	32
4.2	Restoring/improving the hydrological regime	20
6.3	Legal protection of habitats and species	18
4.0	Other wetland-related measures	6
7.1	Regulation/ Management of hunting and taking	4
9.2	Regulating/Managing exploitation of natural resources on sea	4
2.0	Other agriculture-related measures	2
2.1	Maintaining grasslands and other open habitats	2
3.0	Other forestry-related measures	2
4.1	Restoring/improving water quality	2

This information is derived from the Member State national reports submitted to the European Commission under Article 12 of the Birds Directive in 2013 and covering the period 2008-2012. More detailed information, including the MS reports, is available at:

<http://bd.eionet.europa.eu/article12/summary?period=1&subject=A197> .

ⁱ **Assessment of status at the European level:** The EU assessments of birds population status was made by the European Red List of Birds Consortium (under contract with the European Commission)

The EU27 population trends were assessed using these categories: '+' Increasing, '0' Stable, 'F' Fluctuating, '-' Decreasing, 'xu' Uncertain and 'x' Unknown. The breeding population size is estimated in majority of the cases as 'p' number of pairs. Alternative population units used are: 'males' number of males, 'i' number of individuals, 'cmales' number of calling males and 'bfem' number of breeding females. The winter population size is estimated as number of individuals.

ⁱⁱ **Species trends at the Member State level:** The percentage of the EU27 species population occurring in the Member States (% in EU27) is calculated based on the population size reported by the Member States.

ⁱⁱⁱ **Percentage of coverage by Natura 2000 sites:** In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece and for non-Annex I species in the Czech Republic.