European Environment Agency European Topic Centre on Biological Diversity



Riparia riparia

Annex I No International action plan No

Sand Martin, *Riparia riparia*, is a species of passerine bird in the swallow family found in wetland, urban and river and lake ecosystems.


Riparia riparia has a breeding population size of 1260000-2810000 pairs and a breeding range size of 2290000 square kilometres in the EU27. The breeding population trend in the EU27 is Stable in the short term and Decreasing in the long term.

The EU population status of *Riparia riparia* was assessed as Depleted, because the EU27 population or range declined by at least 20% since 1980, but has no longer been declining since 2001.

Assessment of status at the European level

Breeding population size	Breeding population trend		Range _	Breeding range trend		Winter	Winter population trend		_Population
	Short term	Long term	area	Short term	Long term	population size	Short term	Long term	status
1260000 - 2810000 p	0	_	2290000						Depleted


See the endnotes for more informationⁱ




Trends at the Member State level

MS/Ter. % in			Breeding population trend		Range	Breeding range trend		Winter population	Winter population trend	
MS/Tel. EU27	EU27	population size	Short term	Long term	area	Short term	Long term	size	Short term	Long term
AT	1.2	6000 - 9000 p	-	X	25788	0	-			
BE	0.8	5000 - 10000 p	0	0	9949	-	-			
BG	1.5	20000 - 40000 p	0	0	59600	0	0			
CZ										
DE	12.3	105000 - 165000 p	-	0	206002	0	0			
DK	1.8	11000 - 11000 p	-	-	40934	0	0			
EE	1.6	5000 - 10000 p	-	-	40800	0	-			
ES	10.4	540000 - 750000 p	0	-	310731	0	-			
FI	11.5	46000 - 63000 p	-	-	324700	X	-			
FR	8.5	50000 - 100000 p	X	-	273300	-	+			
GR										
HU	3.2	15000 - 50000 p	-	-	34312	-	X			
IE	6.2	144100 - 1113380 i	+	X	82900	+	0			
IT	3.7	6000 - 8000 p	-	-	76500	-	_			
LT	4.9	30000 - 70000 p	-	-	72200	0	0			
LU		110 - 150 p	F	-	1216	F	-			
LV	2.6	11340 - 19920 p	X	-	62329	0	0			
NL	2.2	25000 - 38000 p	+	+	39184	0	0			
PL		150000 - 300000 p	X	X		X	X			
PT	3.6	10000 - 50000 p	X	X	81100	0	+			
RO	0.3	45000 - 90000 p	X	X	101300	X	X			
SE	4.9	26000 - 40000 p	-	-	226700	0	X			
SI	0.2	500 - 2000 p	F	X	1998	+	+			
SK	1.6	5000 - 10000 p	-	-	33697	0	0			
UK	16.9	54000 - 174000 p	+	+	187300	+	-			

See the endnotes for more information $^{\mathrm{ii}}$





Short-term winter population trend was not reported for this species.

Main pressures and threats reported by Member States

For the bird species triggering SPA classification Member States were asked to report the 20 most important pressures and threats using an agreed hierarchical list which can be found on the Article 12 Reference Portal (http://bd.eionet.europa.eu/activities/Reporting/Article_12/reference_portal). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. The table below only contains information from Member States, where a species triggers SPA classification. Pressures and threats were ranked in three classes 'high, medium and low importance', the table below only shows pressures and threats classed as 'high', for some species there were less than ten pressures and threats reported as highly important.

Ten most frequently reported 'highly important' pressures and threats

Code	Activity	Frequency
J02	Changes in water bodies conditions	43
J03	Other changes to ecosystems	29
G01	Outdoor sports, leisure and recreational activities	14
M01	Abiotic changes (climate change)	14

Proportion of population covered by the Natura 2000 network

For the bird species triggering SPA classification Member States were asked to report the size of a species population occurring within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size.

Percentage of coverage by Natura 2000 sites

MS/territory	season	SPA trigger	% coverage
AT	breeding	NO	
BE	breeding	NO	
BG	breeding	YES	18.03
DE	breeding	NO	
DK	breeding	NO	
EE	breeding	NO	
ES	breeding	YES	0.07
FI	breeding	NO	
FR	breeding	NO	
HU	breeding	YES	22.08
IE	breeding	NO	

MS/territory	season	SPA trigger	% coverage
IT	breeding	NO	_
LT	breeding	NO	
LU	breeding	YES	44.04
LV	breeding	NO	
NL	breeding	YES	17.69
PL	breeding	NO	
PT	breeding	NO	
RO	breeding	NO	
SE	breeding	NO	
SI	breeding	NO	
SK	breeding	YES	25.1
UK	breeding	NO	

See the endnotes for more information iii

Most frequently reported conservation measures

For the bird species triggering SPA classification Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 12 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.3	Legal protection of habitats and species	33
6.1	Establish protected areas/sites	27
7.4	Specific single species or species group management measures	13
4.1	Restoring/improving water quality	7
4.2	Restoring/improving the hydrological regime	7
4.4	Restoring coastal areas	7
9.1	Regulating/Management exploitation of natural resources on land	7

This information is derived from the Member State national reports submitted to the European Commission under Article 12 of the Birds Directive in 2013 and covering the period 2008-2012. More detailed information, including the MS reports, is available at: http://bd.eionet.europa.eu/article12/summary?period=1&subject=A249.

Riparia riparia

Report under the Article 12 of the Birds Directive

¹ Assessment of status at the European level: The EU assessments of birds population status was made by the European Red List of Birds Consortium (under contract with the European Commission)

The EU27 population trends were assessed using these categories: '+' Increasing, '0' Stable, 'F' Fluctuating, '-' Decreasing, 'xu' Uncertain and 'x' Unknown. The breeding population size is estimated in majority of the cases as 'p' number of pairs. Alternative population units used are: 'males' number of males, 'i' number of individuals, 'cmales' number of calling males and 'bfem' number of breeding females. The winter population size is estimated as number of individuals.

ⁱⁱSpecies trends at the Member State level: The percentage of the EU27 species population occurring in the Member States (% in EU27) is calculated based on the population size reported by the Member States.

Percentage of coverage by Natura 2000 sites: In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece and for non-Annex I species in the Czech Republic.