

A1.1xx Invertebrate-dominated moderately exposed Pontic mediolittoral rock

Summary

The habitat is present in the Black Sea on areas of moderately exposed mediolittoral rock. It is not present in the Sea of Marmara. Eutrophication is the most significant historical pressure on the habitat. It is also sensitive to coastal development and chemical pollution. The habitat can benefit from future conservation measures including: measures to maintain physical and biological integrity, including pollution control and regulation; Improvement of water quality management outside EU member states; development controls and contingency plans to be followed in the event of a major pollution incident; survey and monitoring programmes and raised public awareness of ecological value and vulnerability.

Synthesis

Detailed information on the abundance and extent of this habitat is lacking. Information on the quantity and quality of this habitat including historical or recent trends is unknown. For the purposes of Red List assessment this habitat is considered to be Data Deficient.

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Data Deficient	-	Data Deficient	-

Sub-habitat types that may require further examination

None

Habitat Type

Code and name

A1.1xx Invertebrate-dominated moderately exposed Pontic mediolittoral rock

No characteristic photograph of this habitat currently available.

Habitat description

These communities are located in the moderately exposed mediolittoral rock zone. They result from a combination of moderate waves, variations in atmospheric pressure, and variations in wind.

This habitat is comprised of all invertebrate communities found in the moderately exposed mediolittoral rock zone.

Indicators of quality:

Both biotic and abiotic indicators have been used to describe marine habitat quality. These include; the presence of characteristic species and those which are sensitive to the pressures the habitat may face, water quality parameters, levels of exposure to particular pressure as well as and more integrated indices which describe habitat structure and function, such as trophic index, or successional stages of development in habitats that have a natural cycle of change over time. There are no commonly agreed indicators of quality for this habitat, although particular parameters may have been set in certain situations e.g. protected features within Natura 2000 sites, where reference values have been determined and applied on a location-specific basis.

Characteristic species:

Mytilids (*Mytilus galloprovincialis*, *Mytilaster lineatus*), barnacles (*Chthamalus stellatus*, *Amphibalanus improvisus*), *Actinia equina*, bryozoans, crust sponges and articulated corallines (*Corallina elongata*, *C. officinalis*).

Classification

This habitat may be equivalent to, or broader than, or narrower than the habitats or ecosystems in the following typologies.

EUNIS (v1405):

Level 4. A sub-habitat of 'Pontic littoral rock' (A1.1).

Annex 1:

1160 Large shallow inlets and bays

1170 Reefs

8330 Submerged or partially submerged sea caves

MAES:

Marine - Marine inlets and transitional waters

Marine - Coastal

MSFD:

Littoral rock and biogenic reef

EUSEaMap:

Not mapped

IUCN:

12.1 Rocky shoreline

Does the habitat type present an outstanding example of typical characteristics of one or more biogeographic regions?

Unknown

Justification

There is insufficient knowledge and information on this habitat to state whether it is an outstanding example of this biogeographic region.

Geographic occurrence and trends


Region	Present or Presence Uncertain	Current area of habitat	Recent trend in quantity (last 50 yrs)	Recent trend in quality (last 50 yrs)
--------	-------------------------------	-------------------------	--	---------------------------------------

Region	Present or Presence Uncertain	Current area of habitat	Recent trend in quantity (last 50 yrs)	Recent trend in quality (last 50 yrs)
<i>Black Sea</i>	Black Sea: Present Sea of Marmara: Present	Unknown Km ²	Unknown	Unknown

Extent of Occurrence, Area of Occupancy and habitat area

	Extent of Occurrence (EOO)	Area of Occupancy (AOO)	Current estimated Total Area	Comment
<i>EU 28</i>	Unknown Km ²	Unknown	Unknown Km ²	The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO.
<i>EU 28+</i>	Unknown Km ²	Unknown	Unknown Km ²	The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO.

Distribution map



There is insufficient data to produce a map of the distribution of this habitat.

How much of the current distribution of the habitat type lies within the EU 28?

It is unknown how much of this habitat is hosted by the EU28 in the Black Sea

Trends in quantity

There is insufficient data to accurately assess changes in quantity of the habitat

- Average current trend in quantity (extent)

EU 28: Unknown

EU 28+: Unknown

- Does the habitat type have a small natural range following regression?

Unknown

Justification

The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO. There is insufficient data to accurately assess whether the habitat has undergone a significant decline (>25% of extent) in the last 50 years.

- Does the habitat have a small natural range by reason of its intrinsically restricted area?

Unknown

Justification

There is insufficient data and knowledge on this habitat to state whether it has a small natural range by reason of an intrinsically restricted area.

Trends in quality

There is insufficient data to accurately assess changes in quality of the habitat

- Average current trend in quality

EU 28: Unknown

EU 28+: Unknown

Pressures and threats

Eutrophication as a result of nutrient enrichment (N, P and organic matter) is the most significant historic pressure on the habitat. Anoxic and hypoxic conditions due to eutrophication caused mass mortalities in faunal communities. Since the 1990s this pressure has reduced due to tighter controls on pollution in the catchment of the Danube and other rivers which enter the north-west Black Sea. Whilst this pressure is now reduced it is still a continuing threat in the current and future periods. This is especially true for non EU countries surrounding the Black Sea which are not bound by the agreements such as the Water Framework Directive (WFD).

The habitat is also likely to be sensitive to:

- Coastal developments including the construction of marinas and slipways, sediment extraction, the widening and dredging of channels, creation of artificial beaches, road developments and sea defenses. These activities may alter the hydrological regime which will in turn affect the character and viability of the habitat.

- Chemical pollution. This is a threat of current and future importance which at its most severe can result in species can lead to mortality. High mortality rates can lead to a reduction in extent. Lower mortality rates will result in a reduction in habitat quality. Chemical pollution may also affect the size and growth rate of some of the associated fauna.

List of pressures and threats

Urbanisation, residential and commercial development

Other urbanisation, industrial and similar activities

Pollution

Nutrient enrichment (N, P, organic matter)

Input of contaminants (synthetic substances, non-synthetic substances, radionuclides) - diffuse sources, point sources, acute events

Conservation and management

Conservation and management measures which would benefit this habitat include implementing measures to maintain physical and biological integrity, including pollution control and regulation, improvement of water quality management outside EU member states, development of controls and contingency plans to be followed in the event of a major pollution incident, survey and monitoring programmes, raised public awareness of ecological value and vulnerability.

List of conservation and management needs

Measures related to marine habitats

Other marine-related measures

Measures related to urban areas, industry, energy and transport

Other measures

Conservation status

Annex 1:

1160: MBL5 U1

1170: MBL5 U1

When severely damaged, does the habitat retain the capacity to recover its typical character and functionality?

There is insufficient data and knowledge of this habitat to assess its capacity to recover

Effort required

10 years
Unknown

Red List Assessment

Criterion A: Reduction in quantity

Criterion A	A1	A2a	A2b	A3
EU 28	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %

There is insufficient data on changes in quantity of this habitat to undertake an assessment using criterion A.

Criterion B: Restricted geographic distribution

Criterion B	B1				B2				B3
	EOO	a	b	c	AOO	a	b	c	
EU 28	unknown Km ²	Unknown							
EU 28+	unknown Km ²	Unknown							

The precise extent of the habitat is unknown. Therefore there is insufficient data to produce EOO and AOO figures.

Criterion C and D: Reduction in abiotic and/or biotic quality

Criteria C/D	C/D1		C/D2		C/D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %

Criterion C	C1		C2		C3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %

Criterion D	D1		D2		D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%
EU 28+	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%

Experts consider there to be insufficient data to conduct an assessment using criteria C/D.

Criterion E: Quantitative analysis to evaluate risk of habitat collapse

Criterion E	Probability of collapse
EU 28	unknown
EU 28+	unknown

There is no quantitative analysis available to estimate the probability of collapse of this habitat type.

Overall assessment "Balance sheet" for EU 28 and EU 28+

	A1	A2a	A2b	A3	B1	B2	B3	C/D1	C/D2	C/D3	C1	C2	C3	D1	D2	D3	E
EU28	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD
EU28+	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Data Deficient	-	Data Deficient	-

Confidence in the assessment

Low (mainly based on uncertain or indirect information, inferred and suspected data values, and/or limited expert knowledge)

Assessors

S. Beal, G. Komakhidze, D. Micu, V. Mihneva, N. Milchakova, B. Yokes

Contributors

S. Beal, G. Komakhidze, D. Micu, V. Mihneva, N. Milchakova, B. Yokes

Reviewers

L. Bat

Date of assessment

19/03/2015

Date of review

20/01/2016

References

- Anon. 2006. *The northwestern part of the Black Sea: biology and ecology*. Kiev: Naukova Dumka. 701pp.
- Bacescu, M. C., Muller G. I., Gomoiu, M-T. 1971.. Cercetari de ecologie bentica in Marea Neagra (analiza cantitativa, calitativa si comparata a faunei bentice pontice). *Ecologie Marina* vol. IV. Editura Academiei R.S.R., Bucuresti, 357 pp.
- Bacescu M., 1977. Les biocenoses benthiques de la Mer Noire. *Biologie des eaux saumâtres de la Mer Noire, Première partie*: 128-134.
- Çulha, M. & Bat, L. 2010. Visible decline of limpet *Patella caerulea* Linnaeus, 1758, a biomonitor species, at the Sinop Peninsula and vicinity (the southern Black sea, Turkey). *Journal of Environmental Protection and Ecology* 11(3): 1024-1029.
- Dimitrova-Konaklieva, S. 2000. *Flora of the Marine Algae of Bulgaria (Rhodophyta, Phaeophyta, Chlorophyta)*. Pensoft, Sofia, Bulgaria.
- Kalugina-Gutnik A.A. 1970. *The composition and distribution of benthic vegetation in the south-eastern part of the Black Sea*. Ecological and morphological studies of benthic organisms. Kiev: Naukova Dumka, p. 185- 202.
- Kalugina-Gutnik A.A. 1975. *Phytobenthos of Black Sea*, Kiev: Naukova Dumka, 275 p.
- Konsulov, A. 1998. *Black Sea Biological Diversity: Bulgaria. Volume 5 of Black Sea environmental series*. United Nations Publications, New York, USA.
- Kopiy, V. G. Bondarenko, L. V. 2012. The community of the macrozoobenthos of mediolittoral zone of Western Crimea. Biodiversity and sustainable development: Abstracts of the II Intern. *scientific and practic Conf., Simferopol*: 189-192.
- Kostenko, N. S. 2003. Some trends of the succesion of bottom vegetation in the Karadag area. *Proc. Sciences. Rec. NaUKMA, Ser. "Biologiya and ekologiya"*: 429-432.
- Marinov, T. 1990. *The zoobenthos from the Bulgarian Sector of the Black Sea*. Publishing house of the Bulgarian Academy of Sciences, Sofia, pp 195 (in Bulgarian).
- Micu, D., Zaharia, T., Todorova, V., Niță, V. 2007. *Romanian Marine Habitats of European Interest*. Punct Ochit Publishers, Constanța, Romania.
- Micu, D. 2008. Open Sea and Tidal Areas. In: Gafta D. and Mountford J.O. (eds.) *Natura 2000 Habitat Interpretation Manual for Romania*. EU publication no. EuropeAid/121260/D/SV/RO, 101pp. ISBN 978-973- 751-697-8.
- Micu, D., Zaharia, T., Todorova, V. 2008. Natura 2000 habitat types from the Romanian Black Sea. In: Zaharia T, Micu D, Todorova V, Maximov V, Niță V. *The development of an indicative ecologically coherent network of marine protected areas in Romania*. Romart Design Publishing, Constanta, Romania.
- Moncheva. S., Todorova, V., (eds). 2013. *Initial assessment of the marine environment*. Article 8,

MSFD 2008/56/EC and NOOSMV (2010). 500p

Petranu, A. 1997. *Black Sea Biological Diversity: Romania. Volume 4 of the Black Sea Environmental Series*. United Nations Publications, New York, USA.

Prodanov, B., Kotsev, I., Keremedichiev, S., Todorova, V., Dimitrov, L. 2013. *Initial assessment of the technogenic pressure in the mediolittoral zone of the Bulgarian black sea coast*. Second European SCGIS Conference "Conservation of Natural and Cultural Heritage for Sustainable Development: GIS-Based Approach", 2013: 4-13.

Salomidi, M., Katsanevakis, S., Damalas, D., Mifsud, R., Todorova, V., Pipitone, C., Fernandez, T. V., Mirto, S., Galparsoro, I., Pascual, M., Borja, Á., Rabaut, M., Braeckman, U. 2010. Monitoring and Evaluation of Spatially Managed Areas. Catalogue of European seabed biotopes. Deliverable 1.2. Available at: <http://www.mesma.org/default.asp?ZNT=S0T1O-1P24>. (Accessed: 19/08/2015).

Terentyev, A. S. 2011. Macrozoobenthos of coastal part of the Kerch Bay (summer, 2009). Ecology of cities and recreational areas. / All_Ukrainian Scientific Conference Proceedings of articles: 261-263.

Todorova, V., Panayotova, M. 2011. *Black mussels and/or barnacle communities on mediolittoral rocks*. Red book of Republic of Bulgaria, Vol. III, Natural habitats, Eds. BAS & MOEW. [ISBN 978-9549746-23-5].

Vershinin, A. 2007. *Life in the Black Sea*. Maccentr, Moscow, Russia.

Zaika V. E., Boltachev A. R., Zuev G. V., Kovalev A. V., Milchakova N. A., Sergeeva N. G. 2004. Floristic and faunistic changes in the Crimean Black Sea shelf after 1995 - 1998, Marine Ecological Journal, 3(2), p. 37-44.

Zaitsev, Y. P., Alexandrov, B. G. 1998. *Black Sea Biological Diversity: Ukraine. Volume 7 of the Black Sea Environmental Series*. United Nations Publications, New York, USA.