

A3.2x Mytilid-dominated Pontic moderately exposed upper infralittoral rock, blocks and boulders, with foliose algae (other than Fucales)

Summary

The habitat is present in the Black Sea on moderately exposed upper infralittoral rock. It is not present in the Sea of Marmara. Eutrophication is the main historic pressure on this habitat. Additional pressures include: coastal development and chemical pollution. Conservation and management measures relevant to this habitat include: measures to maintain physical and biological integrity, improvement of water quality, coastal development controls, contingency plans in case of pollution events and raised public awareness.

Synthesis

Detailed information on the abundance and extent of this habitat is lacking. Information on the quantity and quality of this habitat including historical or recent trends is unknown. For the purposes of Red List assessment this habitat is considered to be Data Deficient.

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Data Deficient	-	Data Deficient	-

Sub-habitat types that may require further examination

None

Habitat Type

Code and name

A3.2x Mytilid-dominated Pontic moderately exposed upper infralittoral rock, blocks and boulders, with foliose algae (other than Fucales)

No photographs are currently available of this habitat.

Habitat description

Rocky habitats in the infralittoral zone exposed to wave action. Rocks are typically covered by foliose algal communities. Mytilids are a constant component. These habitats occur on exposed rocky coasts, from low water up to depths of 10m. Winter storms may cause changes to species compositions and coverage.

Indicators of quality:

A set of categories common to the habitat indicating good biotic and abiotic quality. These indicators will be used, together with the general "Stages of Quality Decline" (applicable for all terrestrial and marine habitats), to assess the severity of quality decline over a certain time frame on a country/regional scale and for EU28 and EU28+.

Characteristic species:

Characteristic species includes: *Mytilus galloprovincialis*, *Mytilaster lineatus*, *Lophosiphonia obscura*, *Ceramium ciliatum*, *Padina pavonica*, *Grateloupia dichotoma*, *Dilophus fasciola*, *F. repens*, *Polysiphonia opaca*, *Ceramium ciliatum*, *Ulva compressa*, *Ulva intestinalis*, *Ulva linza*.

Classification

This habitat may be equivalent to, or broader than, or narrower than the habitats or ecosystems in the following typologies.

EUNIS (2004):

Level 4. A sub-habitat of 'Pontic infralittoral rock' (A3.1)

Annex 1:

1130 Estuaries

1160 Large swallow inlets and bays

1170 Reefs

8330 Submerged or partially submerged sea caves

MAES:

Marine - Marine inlets and transitional waters

Marine - Coastal

MSFD:

Shallow sublittoral rock and biogenic reef

EUSEaMap:

Shallow photic rock or biogenic reef

Shallow aphotic rock or biogenic reef

IUCN:

9.2 Subtidal rock and rocky reefs

Does the habitat type present an outstanding example of typical characteristics of one or more biogeographic regions?

Unknown

Justification

There is insufficient knowledge and information on this habitat to state whether it is an outstanding example of this biogeographic region.


Geographic occurrence and trends

Region	Present or Presence Uncertain	Current area of habitat	Recent trend in quantity (last 50 yrs)	Recent trend in quality (last 50 yrs)
<i>Black Sea</i>	Black Sea: Present Sea of Marmara: Uncertain	Unknown Km ²	Unknown	Unknown

Extent of Occurrence, Area of Occupancy and habitat area

	Extent of Occurrence (EOO)	Area of Occupancy (AOO)	Current estimated Total Area	Comment
EU 28	Unknown Km ²	Unknown	Unknown Km ²	The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO.
EU 28+	Unknown Km ²	Unknown	Unknown Km ²	The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO.

Distribution map



There is insufficient data to produce a map of the distribution of this habitat.

How much of the current distribution of the habitat type lies within the EU 28?

It is unknown how much of this habitat is hosted by the EU28 in the Black Sea.

Trends in quantity

There is insufficient data to accurately assess changes in quantity of the habitat

- Average current trend in quantity (extent)
EU 28: Unknown
EU 28+: Unknown
- Does the habitat type have a small natural range following regression?
Unknown

Justification

The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO

and AOO. There is insufficient data to accurately assess whether the habitat has undergone a significant decline (>25% of extent) in the last 50 years.

- Does the habitat have a small natural range by reason of its intrinsically restricted area?

Unknown

Justification

There is insufficient data and knowledge on this habitat to state whether it has a small natural range by reason of an intrinsically restricted area.

Trends in quality

There is insufficient data to accurately assess changes in quality of the habitat

- Average current trend in quality

EU 28: Unknown

EU 28+: Unknown

Pressures and threats

Eutrophication as a result of nutrient enrichment (N, P and organic matter) is the most significant historic pressure on the habitat. Anoxic and hypoxic conditions due to eutrophication caused mass mortalities in benthic communities. Since the 1990s this pressure has reduced due to tighter controls on pollution in the catchment of the Danube and other rivers which enter the north-west Black Sea. Whilst this pressure is now reduced it is still a continuing threat in the current and future periods. This is especially true for non EU countries surrounding the Black Sea which are not bound by the agreements such as the Water Framework Directive (WFD).

Coastal developments including the construction of marinas and slipways, sediment extraction, the widening and dredging of channels, creation of artificial beaches, road developments and sea defences. These activities may alter the hydrological regime which will in turn affect the character and viability of the habitat.

Chemical pollution. This is a threat of current and future importance which at its most severe can result in species can lead to mortality. High mortality rates can lead to a reduction in extent. Lower mortality rates will result in a reduction in habitat quality. Chemical pollution may also affect the size and growth rate of some of the associated fauna.

List of pressures and threats

Urbanisation, residential and commercial development

Other urbanisation, industrial and similar activities

Pollution

Nutrient enrichment (N, P, organic matter)

Input of contaminants (synthetic substances, non-synthetic substances, radionuclides) - diffuse sources, point sources, acute events

Conservation and management

Conservation and management measures which would benefit this habitat include implementing measures to maintain physical and biological integrity, including pollution control and regulation, improvement of water quality management outside EU member states, coastal development controls, contingency plans to be followed in the event of a major pollution incident, raised public awareness of ecological value and vulnerability.

List of conservation and management needs

Measures related to marine habitats

Other marine-related measures

Measures related to spatial planning

Other spatial measures

Establish protected areas/sites

Measures related to urban areas, industry, energy and transport

Other measures

Conservation status

Annex 1:

1160: MBLS U1

1170: MBLS U1

When severely damaged, does the habitat retain the capacity to recover its typical character and functionality?

There is insufficient data and knowledge of this habitat to assess its capacity to recover

Effort required

10 years
Unknown

Red List Assessment

Criterion A: Reduction in quantity

Criterion A	A1	A2a	A2b	A3
EU 28	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %

There is insufficient data on changes in quantity of this habitat to undertake an assessment using criterion A.

Criterion B: Restricted geographic distribution

Criterion B	B1				B2				B3
	EOO	a	b	c	AOO	a	b	c	
EU 28	unknown Km ²	Unknown	Unknown	unknown	unknown	Unknown	Unknown	unknown	unknown
EU 28+	unknown Km ²	Unknown	Unknown	unknown	unknown	Unknown	-	unknown	unknown

The precise extent of the habitat is unknown. Therefore there is insufficient data to produce EOO and AOO figures.

Criterion C and D: Reduction in abiotic and/or biotic quality

Criteria C/D	C/D1		C/D2		C/D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %

Criterion C	C1		C2		C3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %

Criterion D	D1		D2		D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%
EU 28+	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%

Experts consider there to be insufficient data to conduct an assessment using criteria C/D.

Criterion E: Quantitative analysis to evaluate risk of habitat collapse

Criterion E	Probability of collapse
EU 28	unknown
EU 28+	unknown

There is no quantitative analysis available to estimate the probability of collapse of this habitat type.

Overall assessment "Balance sheet" for EU 28 and EU 28+

	A1	A2a	A2b	A3	B1	B2	B3	C/D1	C/D2	C/D3	C1	C2	C3	D1	D2	D3	E
EU28	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD
EU28+	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Data Deficient	-	Data Deficient	-

Confidence in the assessment

Low (mainly based on uncertain or indirect information, inferred and suspected data values, and/or limited expert knowledge)

Assessors

S. Beal, G. Komakhidze, D. Micu, V. Mihneva, N. Milchakova, B. Yokes

Contributors

S. Beal, G. Komakhidze, D. Micu, V. Mihneva, N. Milchakova, B. Yokes

Reviewers

N. Dankers

Date of assessment

19/03/2015

Date of review

17/12/2015

References

- Afanasiev D. F., Korpakova I. G. 2008. Macrophytobenthos Russian Azov-Black Sea., Rostov-on-Don: FGUP AzNIIRH.
- Anon. 2006. *The northwestern part of the Black Sea: biology and ecology*. Kiev: Naukova Dumka. 701pp.
- Bacescu, M. C., Muller G. I., Gomoiu, M-T. 1971. . Cercetari de ecologie bentica in Marea Neagra (analiza cantitativa, calitativa si comparata a faunei bentice pontice). *Ecologie Marina* vol. IV. Editura Academiei R.S.R., Bucuresti, 357 pp..
- Bacescu M., 1977. Les biocenoses benthiques de la Mer Noire. *Biologie des eaux saumâtres de la Mer Noire, Premiere partie*: 128-134.
- Chernyakov D. A. 1995. Natural-aquatic landscape complexes of the Tendra and Egorlyk bays and monitoring of their state in Black Sea Biosphere Reserve.
- Çulha, M., Bat, L., Türk Çulha, S. & Çelik, M.Y. 2010. Benthic mollusk composition of some facies in the upper-infralittoral zone of the southern Black Sea, Turkey. *Turkish Journal of Zoology* 34: 523-532.
- Dimitrova-Konaklieva, S. 2000. *Flora of the Marine Algae of Bulgaria (Rhodophyta, Phaeophyta, Chlorophyta)*. Pensoft, Sofia, Bulgaria.
- Kalugina-Gutnik A.A. 1970. *The composition and distribution of benthic vegetation in the south-eastern part of the Black Sea*. Ecological and morphological studies of benthic organisms. Kiev: Naukova Dumka, p. 185- 202.
- Kalugina-Gutnik A.A. 1975. *Phytobenthos of Black Sea*, Kiev: Naukova Dumka, 275 p. Kiseleva, M. I. 1981. *Benthos of Black Sea mobile substrates*. Naukova dumka, Kiev, pp 165.
- Konsulov, A. 1998. *Black Sea Biological Diversity: Bulgaria. Volume 5 of Black Sea environmental series*. United Nations Publications, New York, USA.
- Kostenko, N. S. 2003. Some trends of the succession of bottom vegetation in the Karadag area. *Proc. Sciences. Rec. NaUKMA, Ser. "Biologiya and ekologiya"*: 429-432.
- Lisovskaya O.A., Stepanyan O.V. 2009. A variety of coastal macroalgae Taman Peninsula (Russia) in summer., *Algology* 19(4): 341-348.
- Marinov, T. 1990. *The zoobenthos from the Bulgarian Sector of the Black Sea*. Publishing house of the Bulgarian Academy of Sciences, Sofia, pp 195 (in Bulgarian).
- Micu, D., Micu, S. 2006. *Recent records and proposed IUCN status of Donacilla cornea (Poli, 1795) (Bivalvia: Veneroidea: Mesodesmatidae) in the Romanian Black Sea*. *Cercet Mar* 36: 117-132.
- Micu, D., Zaharia, T., Todorova, V., Niță, V. 2007. *Romanian Marine Habitats of European Interest*. Punct Ochit Publishers, Constanța, Romania.
- Micu, D. 2008. Open Sea and Tidal Areas. In: Gafta D. and Mountford J.O. (eds.) *Natura 2000 Habitat Interpretation Manual for Romania*. EU publication no. EuropeAid/121260/D/SV/RO, 101pp. ISBN 978-973- 751-697-8.

- Micu, D., Zaharia, T., Todorova, V. 2008. Natura 2000 habitat types from the Romanian Black Sea. In: Zaharia T, Micu D, Todorova V, Maximov V, Niță V. *The development of an indicative ecologically coherent network of marine protected areas in Romania*. Romart Design Publishing, Constanta, Romania.
- Moncheva. S., Todorova, V., (eds). 2013. *Initial assessment of the marine environment*. Article 8, MSFD 2008/56/EC and NOOSMV (2010). 500p
- Morozova-Vodyanitskaya N. V. 1959. Bottom vegetation of the Black Sea. *Proceedings of the Sevastopol Biological Station* 11: 3 – 28.
- Petranu, A. 1997. *Black Sea Biological Diversity: Romania. Volume 4 of the Black Sea Environmental Series*. United Nations Publications, New York, USA.
- Salomidi, M., Katsanevakis, S., Damalas, D., Mifsud, R., Todorova, V., Pipitone, C., Fernandez, T. V., Mirto, S., Galparsoro, I., Pascual, M., Borja, Á., Rabaut, M., Braeckman, U. 2010. Monitoring and Evaluation of Spatially Managed Areas. Catalogue of European seabed biotopes. Deliverable 1.2. Available at: <http://www.mesma.org/default.asp?ZNT=S0T1O-1P24>. (Accessed: 19/08/2015).
- Teyubova V. F. 2005. Features interannual dynamics species composition and structure macrophytobenthos in the Bay of Novorossiysk (Black sea). *Ekologiya Morya* 69: 53 – 57.
- Teyubova V.F. 2012. *The diversity and ecological features macrophytobenthos the Russian sector of the Black Sea.*, Dissertation on competition degree of candidate of biological sciences, 280 pp.
- Tkachenko, F. P., Kovtun O. O. 2014. Contemporary condition of seaweeds flora of Zmeiny island costal zone (Black Sea). *Chornomors'k. bot. z.* 10(1): 37-47.
- Vershinin, A. 2007. *Life in the Black Sea*. Maccentr, Moscow, Russia.
- Zaika V. E., Boltachev A. R., Zuev G. V., Kovalev A. V., Milchakova N. A., Sergeeva N. G. 2004. Floristic and faunistic changes in the Crimean Black Sea shelf after 1995 – 1998, *Marine Ecological Journal*, 3(2), p. 37-44.
- Zaitsev, Y. P., Alexandrov, B. G. 1998. *Black Sea Biological Diversity: Ukraine. Volume 7 of the Black Sea Environmental Series*. United Nations Publications, New York, USA.